



Otic Anti-Infective/Anesthetic

Therapeutic Class Review (TCR)

October 3, 2011

No part of this publication may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, recording, digital scanning, or via any information storage or retrieval system without the express written consent of Provider Synergies, L.L.C.

All requests for permission should be mailed to:

Attention: Copyright Administrator
Intellectual Property Department
Provider Synergies, L.L.C.
10101 Alliance Road, Suite 201
Cincinnati, Ohio 45242

The materials contained herein represent the opinions of the collective authors and editors and should not be construed to be the official representation of any professional organization or group, any state Pharmacy and Therapeutics committee, any state Medicaid Agency, or any other clinical committee. This material is not intended to be relied upon as medical advice for specific medical cases and nothing contained herein should be relied upon by any patient, medical professional or layperson seeking information about a specific course of treatment for a specific medical condition. All readers of this material are responsible for independently obtaining medical advice and guidance from their own physician and/or other medical professional in regard to the best course of treatment for their specific medical condition. This publication, inclusive of all forms contained herein, is intended to be educational in nature and is intended to be used for informational purposes only. Send comments and suggestions to PSTCEditor@magellanhealth.com.

FDA-APPROVED INDICATIONS^{1,2}

| Drug Name | Manufacturer | Indication(s) |
|--|--|---|
| acetic acid (Vosol®) | generic | For the treatment of otitis externa |
| acetic acid / hydrocortisone (Vosol® HC) | generic | For the treatment of otitis externa |
| acetic acid in aluminum acetate | generic | For the treatment of otitis externa |
| acetic acid/ antipyrine/ benzocaine/ polycosan- ol (Otic Care®, PR Otic®, Treagan®) | Pur Tek, PruGen, Trigen Laboratories | Acute otitis media of various etiologies <ul style="list-style-type: none"> ▪ Prompt relief of pain and reduction of inflammation in the congestive and serous stages. ▪ Adjuvant therapy during systemic antibiotic administration for resolution of the infection. Facilitates the removal of excessive or impacted cerumen |
| antipyrine/ benzocaine | generic | <ul style="list-style-type: none"> ▪ For the treatment of otitis externa ▪ To facilitate the removal of excessive or impacted cerumen |
| antipyrine/ benzocaine/ zinc (Neotic®, Otozin™) | Arbor, Allegis | Acute otitis media of various etiologies <ul style="list-style-type: none"> ▪ Prompt relief of pain and reduction of inflammation in the congestive and serous stages. ▪ Adjuvant therapy during systemic antibiotic administration for resolution of the infection. To facilitate the removal of excessive or impacted cerumen |
| benzocaine (Pinnacaine®) | Larken | For the relief of minor ear pain and itching |
| chloroxylenol / benzocaine / hydrocortisone (Myoxin®, Trioxin®) | GM Pharmaceuticals, Vertical Pharmaceuticals | For the treatment of otitis externa complicated by inflammation and to control pruritus in the auditory canal |
| chloroxylenol/ pramoxine (Pramotic®) | Hawthorn | For the treatment of otitis externa |
| chloroxylenol/ pramoxine/ hydrocortisone (Pramoxine HC®) | | For the treatment of otitis externa |
| chloroxylenol/ pramoxine/ zinc/ glycerin (Zinotic®, Zinotic® ES) | Arbor | For the treatment of otitis externa, including the accompanying pain and itching |

The following products are discontinued – chloroxylenol/pramoxine/ hydrocortisone (Cortamox®), antipyrine/benzocaine/phenylephrine (Chlorphenylcaine®), acetic acid (Borofair®).

OVERVIEW

The standard treatment for acute otitis media (AOM) is the use of systemic antibiotics while topical (otic) therapy antibiotic is recommended for uncomplicated for otitis externa.

Acute otitis externa (AOE) is an acute inflammation of the external auditory canal. Commonly referred to as “swimmer’s ear” or “tropical ear”, this condition is often precipitated by water exposure or trauma. The etiology is typically bacterial; the most common pathogens implicated are *Pseudomonas aeruginosa* and *Staphylococcus aureus*, often occurring as a polymicrobial infection. Patients will

typically complain of otalgia and otorrhea, and the ear canal may appear erythematous and swollen. It is imperative that the ear canal be cleared of any discharge or debris that can occlude the canal since the presence of such material can keep the canal moist and interfere with topical treatment. All ages are affected, with a peak incidence in children aged seven to 12 years.

In 2006, the American Academy of Otolaryngology – Head and Neck Surgery Foundation (AAO-HNSF) released guidelines for the management of acute otitis externa (AOE) in patients over two years of age.³ They recommend the use of topical preparations for initial therapy of diffuse, uncomplicated AOE. Systemic antimicrobial therapy should not be used unless there is extension outside the ear canal or the presence of specific host factors that would indicate a need for systemic therapy (e.g., diabetes, immune deficiency, nonintact tympanic membrane, presence of tympanostomy tubes). Management of AOE should include an assessment of pain and analgesic treatment prescribed based on pain severity. Effective topical treatments include antibiotics (e.g., aminoglycosides, polymyxin B, quinolones), steroids (e.g., hydrocortisone), and low pH antiseptics, (e.g., aluminum acetate or acetic acid).⁴

Evidence supports equivalent results with ear cleansing, an ear wick, and any of the choices of topical agents including acidifying agents, antibiotics, antibiotic and steroid combinations, or antifungal agents.⁵ The choice of therapy should be based on efficacy, low incidence of adverse events, and the likelihood of adherence to therapy.⁶

Cerumen impaction is an accumulation of cerumen that causes symptoms or prevents assessment of the ear canal and/or tympanic membrane.⁷ Cerumen impaction is more common in the elderly. Excessive cerumen normally expelled from the ear canal automatically, assisted by motion of the jaw. The AAO-HNSF clinical practice guidelines for cerumen impaction recommend three options for cerumen removal: topical cerumenolytic agents, irrigation, and manual removal other than irrigation. Cerumenolytics, or wax-softening agents, are used to disperse the cerumen and can be used alone or in combination with irrigation or manual removal. The guidelines state that no specific agent is superior to another and none are found to be superior to either saline or water.

PHARMACOLOGY^{8,9}

Acetic acid is a low pH antiseptic agent. Its mechanism of action is unknown.

Chloroxylenol, a halogenated phenol, provides topical germicidal activity against gram negative and gram positive bacteria, fungi, and yeast.

Hydrocortisone is a corticosteroid that control inflammation, edema, pruritus, and other dermal reactions.

Antipyrine, an anti-inflammatory and analgesic agent, exerts its effects by decreasing prostaglandin synthesis.

Benzocaine and pramoxine are local anesthetics that produce a reversible blockade of nerve conduction by decreasing nerve membrane permeability to sodium, leading to a decreased rate of membrane depolarization and increased threshold for electrical excitability.

Polycosanol is an astringent. It's drying effect and removal of debris helps relieve local burning, itching, and irritation.

Glycerin and zinc acetate are topical skin protectants that may also help provide relief to skin surfaces. Zinc also absorbs moisture, thereby acting as a mild astringent.

PHARMACOKINETICS¹⁰

Due to the topical application of these products, minimal systemic absorption is expected.

CONTRAINDICATIONS/WARNINGS^{11,12}

All products in this review are contraindicated in people with hypersensitivity to any of the ingredients contained in the product, and in persons with perforated tympanic membrane or unexplained discharge from the ear.

These agents should be discontinued if sensitization or irritation occurs.

Pain suppression with the use of benzocaine otic preparations (Pinnacaine, Myoxin, Trioxin) may mask the progression of underlying disease.¹³ Patients prescribed a topical anesthetic should be re-examined within 48 hours to ensure appropriate response to primary therapy.

Overgrowth of non-susceptible organisms or superinfection may occur with prolonged use of products containing a germicidal agent, such as acetic acid (Vosol, Vosol HC, Otic Care, PR Otic, Treagan) or chloroxylonol (Myoxin, Trioxin, Pramotic, Pramoxine HC, Zinotic, Zinotic ES).

Studies evaluating cerumenolytics excluded patients with otitis externa; therefore, cerumenolytics should be avoided in patients with active infections of the ear canal.¹⁴

These agents are for otic use only. Avoid accidental exposure to the eyes.

DRUG INTERACTIONS¹⁵

No clinically relevant drug-drug interactions are expected with agents in this review when used as an otic agent.

ADVERSE EFFECTS^{16,17}

The products in this review are generally tolerated well; however, stinging, burning local irritation, contact dermatitis, or hypersensitivity reactions can occur.

Topical benzocaine may cause contact dermatitis that can worsen or prolong otitis externa.

Discontinue treatment if signs and symptoms (e.g., burning, pruritus, redness, or oozing sores in ear) of a local allergic reaction occur.

Warming an otic product by holding the bottle in hand for one to two minutes may minimize dizziness and pain on application; do not heat above body temperature.

SPECIAL POPULATIONS^{18,19}

Pediatrics

Children may absorb proportionally larger amounts of topical corticosteroids, such as hydrocortisone (Myoxin, Pramoxine HC, Trioxin, Vosol HC) and therefore may be more susceptible to systemic toxicity, such as HPA axis suppression and Cushing's syndrome. Pediatric patients should be given the least

amount needed for an effective therapeutic regimen. Chronic corticosteroid therapy may interfere with the growth and development of children.

Safety and effectiveness of acetic acid containing products (Vosol, Vosol HC, PR Otic, Treagan) in pediatric patients below the age of three years have not been established.

Do not use benzocaine (Pinnacaine) in infants under one year of age.

The safety and efficacy of chloroxylenol/hydrocortisone/pramoxine otic combinations (Pramoxine HC) has not been established in infants.

Dosing regimens for use in infants and children are available for chloroxylenol/benzocaine/hydrocortisone (Myoxin, Trioxin). Dosing regimens for use in children are available for antipyrine/benzocaine, antipyrine/benzocaine/polycosanol (Otic Care, PR Otic, Treagan), and antipyrine/benzocaine/zinc (Neotic, Otozin).

The prescribing information for chloroxylenol/pramoxine/zinc/glycerin (Zinotic, Zinotic ES) state to use as directed by a physician in infants and children less than two years of age.

Pregnancy

All agents in this review are classified as Pregnancy Category C.

DOSAGE AND ADMINISTRATION^{20,21}

| Drug | Indication/Dosages | Package size |
|--|--|---|
| acetic acid (Vosol) | Instill four to six drops into ear canal(s). Repeat every two to three hours. | 2% - 15 mL |
| acetic acid / hydrocortisone (Vosol HC) | Insert a cotton wick into the ear canal and saturate with medication. Keep wick moist by adding three to five drops every four to six hours. May remove the wick after 24 hours and continue to instill three to five drops into the ear canal(s) three to four times per day. The lower end of the dosage range is usually recommended in children. | 2% /1% - 10 mL |
| acetic acid in aluminum acetate | Instill four to six drops into the ear canal(s). Repeat every two to three hours. | 2% - 60 mL |
| acetic acid/ antipyrine/ benzocaine/ polycosanol (PR Otic, Otic Care, Treagan) | Acute otitis media: Fill ear canal(s) with medication. Repeat every one to two hours until pain and congestion are relieved. Removal of cerumen: Instill in ear canal three times daily for two to three days. | 0.01% / 5.4% / 1.4% / 0.01% - 14 mL, 15 mL |

Dosage and Administration (continued)

| Drug | Indication/Dosages | Package size |
|---|--|---|
| antipyrine/ benzocaine | Otitis externa: Adults: Instill two to four drops into ear canal(s); Repeat three to four times per day, or up to once every one or two hours. Children: Fill ear canal(s) with medication; Repeat every one to two hours as needed for ear pain. Removal of cerumen: Adults, Adolescents and Children: Instill drops into ear canal(s) three to four times per day for two to three days. | 5.4% / 1.4% - 10 mL, 15 mL |
| antipyrine/ benzocaine/ zinc (Neotic, Otozin™) | Acute otitis media: Fill ear canal(s) with medication. Repeat every one to two hours until pain/congestion is relieved. Removal of cerumen: Instill in ear canal(s) three times daily for two to three days. | 5.4%/ 1%/ 1% - 10 mL x 2 |
| benzocaine (Pinnacaine) | Instill four to five drops into ear canal(s). May repeat every one to two hours, as needed. | 20% - 15mL |
| chloroxylenol / benzocaine / hydrocortisone (Myoxin, Trioxin) | Adults and adolescents: Instill four to five drops into ear canal(s) three to four times per day. Maximum of 20 drops per day. Infants and children: Instill three drops into ear canal(s) three to four times per day. For larger children may consider four to five drops three to four times per day. Maximum of 20 drops per day for children, 12 drops per day for infants | 0.1%/ 1.5%/ 1% - 15 mL |
| chloroxylenol/ pramoxine (Pramotic) | Adults: Instil four to five drops into ear canal(s) three to four times per day. Pediatric: Instil three drops into ear canal(s) three to four times per day. | 0.1%/ 1% - 10 mL |
| chloroxylenol/ pramoxine / hydrocortisone (Pramoxine HC) | Instill three to five drops into ear canal(s) three to four times per day. | 0.1%/ 1%/ 1% - 10 mL |
| chloroxylenol/ pramoxine/ zinc/ glycerin (Zinotic, Zinotic ES) | Adults and adolescents: Instill four to five drops into ear canal(s) three to four times per day for up to 10 days. Maximum 20 drops per ear per day. Children ≥ two years: Instill three drops into ear canal(s) three to four times per day for up to 10 days. Maximum 20 drops per ear per day. Infants and children < two years of age: use as directed by physician. Maximum 12 drops per ear per day for children < two years; no recommendation is available for infants. | Zinotic – 0.1%/ 0.5%/ 0.1%/ 0.1% - 15 mL Zinotic ES – 0.1%/ 1%/ 1%/ 1% - 15 mL |

Patients should be instructed to remain still with while medication is in the ear canal for three to five minutes.²²

If obstruction in the ear canal is evident, aural cleansing (aural toilet) to remove obstructing debris and placement of a wick if edema prevents drug delivery should be considered.²³

In general, it is recommended that drops be given for three days beyond the cessation of symptoms (typically five to seven days); however, in patients with more severe infections, 10 to 14 days of treatment may be required.²⁴

CLINICAL TRIALS

Search Strategy

Search strategy included the use of all drugs in this class and the FDA-approved indications performed on PubMed and included studies for analysis in the review published in English, performed with human participants and where allocation was randomized and a control or comparison group was included allocated participants to comparison groups. Limited comparative clinical trials were found.

Acetic acid versus triamcinolone/acetic acid versus dexamethasone/neomycin/polymyxin

In a double-blind trial 213 adults with acute otitis externa were randomized to receive acetic acid, 0.1% triamcinolone acetate and acetic acid, or 0.66 mg dexamethasone phosphate sodium, 5 mg neomycin sulphate, and 10,000 IU polymyxin B sulphate/mL.²⁵ Each patient received three drops of the study preparation three times daily. The primary outcome measure was the duration of symptoms in days until recovery according to the patient diary entries. Secondary outcome measures were the cure rate at days six-eight, 13-15, and 20-22 and the recurrence of symptoms between days 21 and 42. The median duration to recovery differed between the treatment arms: 8 days (95% CI; 7 to 9) in the acetic acid group, 7 days (5.8 to 8.3) in the steroid and acetic acid group, and 6 days (5.1 to 6.9) in the steroid and antibiotic group. The overall cure rate for all treatment groups after 7, 14, and 21 days was 40, 72, and 79 percent, respectively. Otoscopic signs were not significantly different between treatment arms. Significantly more patients in the acetic acid group had recurrence of symptoms between days 21 and 42 and more severe symptoms than in the other two groups.

Aluminum acetate versus gentamicin

The clinical efficacy of the topical antiseptic, aluminium acetate, and the topical antibiotic, gentamicin was compared for the initial treatment of otorrhea.²⁶ The study included 139 affected ears of which 102 (74 percent) completed the trial. There was no significant difference between the two treatment groups in improvement in otorrhea (67 percent for aluminum acetate versus 68 percent for gentamicin). No resistant organisms to aluminium acetate were reported; however twelve gentamicin-treated ears had gentamicin resistant organisms and one patient developed a gentamicin-resistant *Pseudomonas*.

SUMMARY

The products in this review contain various combinations of antiseptic, anesthetic, astringent, anti-inflammatory and analgesic compounds and are used topically for the treatment of otitis externa, as adjunct therapy in the treatment of acute otitis media, and to facilitate cerumen removal from the ear canal.

The American Academy of Otolaryngology – Head and Neck Surgery Foundation (AAO-HNSF) recommends topical preparations for initial therapy of uncomplicated acute otitis externa; while systemic antimicrobial therapy should be used if there is extension outside the ear canal or if specific host factors are present.

Similar results have been reported with any of the choices of topical agents including acidifying agents, antibiotics, antibiotic and steroid combinations, or antifungal agents in the treatment of otitis externa. In addition, for cerumen removal, no specific agent was found to be superior to another and none were superior to either saline or water.

All products in this review are contraindicated in persons with perforated tympanic membranes or unexplained discharge from the ear.

REFERENCES

- 1 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 2 Drugs.com. Accessed September 29, 2011.
- 3 Rosenfeld RM, Brown L, Cannon CR, et al for the American Academy of Otolaryngology--Head and Neck Surgery Foundation. Clinical practice guideline: acute otitis externa. Otolaryngol Head Neck Surg. 2006; 134(4 Suppl):S4-23. Available at: <http://www.entnet.org/Practice/clinicalPracticeguidelines.cfm>. Accessed September 29, 2011.
- ⁴ Wan-Chih. Drug Treatments for Swimmer's Ear: An Update. Pharmacist Letter. 2009; 25(8):250808.
- 5 Holten KB, Glick J. Management of the Patient with Otitis Externa. Journal of Family Practice / April, 2001.
- 6 Rosenfeld RM, Brown L, Cannon CR, et al for the American Academy of Otolaryngology--Head and Neck Surgery Foundation. Clinical practice guideline: acute otitis externa. Otolaryngol Head Neck Surg. 2006; 134(4 Suppl):S4-23. Available at: <http://www.entnet.org/Practice/clinicalPracticeguidelines.cfm>. Accessed September 29, 2011.
- 7 Roland PS, Smith TL, Schwartz SR, et. al. Clinical practice guideline: Cerumen impaction. Otolaryngology--Head and Neck Surgery (2008) 139, S1-S21. Available at: <http://www.entnet.org/Practice/cerumenImpaction.cfm>. Accessed September 29, 2011.
- 8 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 9 Drugs.com. Accessed September 29, 2011.
- 10 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 11 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 12 Drugs.com. Accessed September 29, 2011.
- 13 Rosenfeld RM, Brown L, Cannon CR, et al for the American Academy of Otolaryngology--Head and Neck Surgery Foundation. Clinical practice guideline: acute otitis externa. Otolaryngol Head Neck Surg. 2006; 134(4 Suppl):S4-23. Available at: <http://www.entnet.org/Practice/clinicalPracticeguidelines.cfm>. Accessed September 29, 2011.
- 14 Roland PS, Smith TL, Schwartz SR, et. al. Clinical practice guideline: Cerumen impaction. Otolaryngology--Head and Neck Surgery (2008) 139, S1-S21. Available at: <http://www.entnet.org/Practice/cerumenImpaction.cfm>. Accessed September 29, 2011.
- 15 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 16 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 17 Drugs.com. Accessed September 29, 2011.
- 18 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 19 Drugs.com. Accessed September 29, 2011.
- 20 Clinical Pharmacology. Available at: <http://www.clinicalpharmacology-ip.com/Default.aspx>. Accessed September 29, 2011.
- 21 Drugs.com. Accessed September 29, 2011.
- 22 Rosenfeld RM, Brown L, Cannon CR, et al for the American Academy of Otolaryngology--Head and Neck Surgery Foundation. Clinical practice guideline: acute otitis externa. Otolaryngol Head Neck Surg. 2006; 134(4 Suppl):S4-23. Available at: <http://www.entnet.org/Practice/clinicalPracticeguidelines.cfm>. Accessed September 29, 2011.
- 23 Rosenfeld RM, Brown L, Cannon CR, et al for the American Academy of Otolaryngology--Head and Neck Surgery Foundation. Clinical practice guideline: acute otitis externa. Otolaryngol Head Neck Surg. 2006; 134(4 Suppl):S4-23. Available at: <http://www.entnet.org/Practice/clinicalPracticeguidelines.cfm>. Accessed September 29, 2011.
- 24 Sander R. Otitis Externa: A Practical Guide to Treatment and Prevention. *Am Fam Physician*. 2001 Mar 1;63(5):927-937.
- 25 van Balen FA, Smit WM, et al. Clinical efficacy of three common treatments in acute otitis externa in primary care: randomised controlled trial. *BMJ*. 2003 November 22; 327(7425): 1201-1205.
- 26 Clayton MJ, Osborne JE, Rutherford D, et al. A double-blind, randomized, prospective trial of a topical antiseptic versus a topical antibiotic in the treatment of otorrhoea. *Clin Otolaryngol Allied Sci* 1990;15:7-10.